

LITERATURE CITED

- Bury, R.B. and J.A. Whelan. 1984. Ecology and management of the bullfrog. U.S. Fish and Wildlife Service Resource Publication 155. 23 pp.
- Cook, D. 1997. Microhabitat use and reproductive success of the California red-legged frog (Rana aurora draytonii) and bullfrog (Rana catesbeiana) in an ephemeral marsh. M.S. Thesis, Sonoma State University, California. 47 pp.
- Emlen, S.T. 1977. "Double clutching" and its possible significance in the bullfrog. *Copeia* 1977(4):749-751.
- Fahrig, L.J.H., S.E. Pope, P.D. Taylor, and J.F. Wegner. 1995. Effects of road traffic on amphibian density. *Biological Conservation* 73:177-182.
- Griffiths, R.A. 1997. Temporary ponds as amphibian habitats. *Aquatic Conservation: Marine and Freshwater Ecosystems* 7:119-126.
- Harris, L.D. 1988. The Nature of Cumulative Impacts on Biotic Diversity of Wetland Vertebrates. *Environ Manag* Vol. 12(5):675-693.
- Harvey, H.T. and Associates. (1997). Santa Clara Valley Water District: California red-legged frog distribution and status-1997. Project No. 1164-01.
- Hayes, M.P. and M.R. Jennings. 1988. Habitat correlates of distribution of the California red-legged frog (Rana aurora draytonii) and the foothill yellow-legged frog (Rana boylii): Implications for management. Pages 144-158 In: R. Sarzo, K. E. Severson, and D. R. Patton (technical coordinators). *Proceedings of the Symposium on the Management of Amphibians, Reptiles, and Small Mammals in North America*. U.S.D.A. Forest Service General Technical Report RM-166.
- Hayes, M.P. and M.R. Jennings. 1986. Decline of ranid frog species in western Nrh America: Are bullfrogs (Rana catesbeiana) responsible? *Journal of Herpetology* 20:490-509.
- Hayes, M.P. and D.M. Krempels. 1986. Vocal sac variation among frogs of the genus *Rana* from western North America. *Copeia* 1986(4):927-936.
- Hayes, M.P. and M.R. Tennant. 1985. Diet and feeding behavior of the California red-legged frog, (Rana aurora draytonii) (Ranidae). *The Southwestern Naturalist* 30(4):601-605.
- Hayes, M.P. and M.M. Miyamoto. 1984. Biochemical, behavioral and body size differences between Rana aurora aurora and R. a. draytonii. *Copeia* 1984(4):1018-1022.
- Hitchings, S.P. and T.J.C. Beebee. 1997. Genetic substructuring as a result of barriers to gene flow in urban Rana temporaria (common frog) populations: implications for biodiversity conservation. *Heredity* 79:117-127.

- Jennings, M.R. 1988. Natural History and decline of native ranids in California. Pages 61-72 In: H.F. Delisle, P.R. Brown, B. Kaufman, and B.M. McGuirty (editors). Proceedings of the conference on California herpetology. Southwestern Herpetologists Society, Special Publication (4):1-143.
- Jennings, M.R. and M.P. Hayes. 1985. Pre-1900 overharvest of California red-legged frog (Rana aurora draytonii): The inducement for bullfrog (Rana catesbeiana) introduction. Herpetological Review 31:94-103.
- Jennings, M.R., M.P. Hayes, and D.C. Holland. 1992. A petition to the U.S. Fish and Wildlife Service to place the California red-legged frog (Rana aurora draytonii) and the western pond turtle (Clemmys marmorata) on the list of endangered and threatened wildlife and plants. 21 pp.
- Kruse, K.C. and M.G. Francis. 1977. A predation deterrent in larvae of the bullfrog, Rana catesbeiana. Transactions of the American Fisheries Society 106(3):248-252.
- Laan, R. and B. Verboom. 1990. Effects of pool size and isolation on amphibian communities. Biol. Cons. 54:251-262.
- Lawler, S. P., D. Dritz, T. Strange, and M. Holyoak. 1999. Effects of introduced mosquitofish and bullfrogs on the threatened California red-legged frog. Cons. Biol. 13:613.622.
- Mann, W., P. Dorn, and R. Brandl. 1991. Local distribution of amphibians: the importance of habitat fragmentation. Global Ecology and Biogeography Letters 1:36-41.
- Marsh, D. M., E. H. Fegraus, and S. Harrison. 1999. Effects of breeding pond isolation on the spatial and temporal dynamics of pond use by the tungara frog, Physalaemus pustulosus. J. Anim. Ecol. 68:804-814.
- Mensing, D.M., S.M. Galatowitsch, and J.R. Tester. 1998. Anthropogenic effects on the biodiversity of riparian wetlands of a northern temperate landscape. Journal of Environmental Management 53:349-377.
- Onorata, D., K.R. Marion, and R.A. Angus. 1998. Longitudinal Variations in the Ichtyofaunal assemblages of the upper Cahabe River: possible effects of urbanization in a watershed. Journal of Freshwater Ecology 13:139-154.
- Reh, W. and A. Seitz. 1990. The influence of land use on the genetic structure of populations of the common frog Rana temporaria. Biol Cons. 54:239-249.
- Reinelt, L.E. and B.L. Taylor. 1997. Effects of watershed development on hydrology. Pp. 141-155 In A.L. Azous and R.R. Horner (eds.). Wetlands and Urbanization: Implications for the Future. Final Report of the Puget Sound Wetlands and Stormwater Management Research Program. Available on the Internet at: <http://splash.metrokc.gov/wlr/basins/weturban.htm>.
- Richter, K.O., and A. L. Azous 1997. Amphibian distribution, abundance and habitat use. Pp.

95-110 In A.L. Azous and R.R. Horner (eds.). Wetlands and Urbanization: Implications for the Future. Final Report of the Puget Sound Wetlands and Stormwater Management Research Program. Available on the Internet at:
<http://splash.metrokc.gov/wlr/basins/weturban.htm>.

Richter, K. O. and A. L. Azous. 1995. Amphibian occurrence and wetland characteristics in the Puget Sound Basin. *Wetlands* 15:306-312.

Sjogren-Gulve, P. 1994. Distribution and extinction patterns within a northern metapopulation of the pool frog, (*Rana lessonae*). *Ecology* 75:1357-1367.

Skinner, L., A. de Peyster, and K. Schiff. 1999. Developmental effects of urban storm water in Medaka (*Oryzias latipes*) and inland silverside (*Menidia beryllina*). *Arch. Environ. Contam. Toxicol.* 37:227-235.

Soule, M. E. 1991. Theory and Strategy; Conservation corridors: countering habitat fragmentation. Pp 91-104 In Hudson, W.E., ed. Landscape Linkages and Biodiversity. Washington, DC: Island Press, 1991.

Stebbins, R.C. 1985. A field guide to western reptiles and amphibians. Houghton Mifflin Company, Boston, MA. xiv + 336 pp.

Storer, T.I. 1925. A synopsis of the amphibia of California. University of California Publications in Zoology 27:1-342.

Storer, T.I. 1933. Frogs and their commercial use. *California Fish and Game* 19(3)203-213.

Twedt, B. 1993. A comparative ecology of *Rana aurora* Baird and Girard and *Rana catesbeiana* Shaw at Freshwater Lagoon, Humboldt County, California. Unpubl. M.S. Humboldt State University, Arcata. 53pp + appendix.

USEPA. U.S. Environmental Protection Agency. 1993. Natural Wetlands and Urban Stormwater : Potential Impacts and Management. EPA: Washington DC. Available through EPA Wetlands Hotline. 1-800-832-7828.

U.S. Fish and Wildlife Service. 2000. Draft Recovery Plan for the California red-legged frog (*Rana aurora draytonii*). U.S. Fish and Wildlife Service, Portland, Oregon. 258 pp.

Wright, A.H. and A.A. Wright. 1949. Handbook of frogs and toads of the United States and Canada. Comstock Publishing Company, Inc., Ithaca, NY. xii + 640 pp.